





NARMS - EB 1997

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint

■ Slaughter n=214;


■ Diagnostic n=153

NARMS - EB 1997


Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken


Ceftiofur




Ceftriaxone



Cephalothin



Chloramphenicol







⬇ Breakpoint

■ Slaughter n=214; ■ Diagnostic n=153

NARMS - EB 1997

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint


■ Slaughter n=214;

■ Diagnostic n=153

NARMS - EB 1997

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint


■ Slaughter n=214;

■ Diagnostic n=153

NARMS - EB 1997

Veterinary Isolates

Fig. 4. Minimum Inhibitory Concentrations by Antimicrobial Agent for *Salmonella* Isolates from Chicken



⬇ Breakpoint

■ Slaughter n=214; ■ Diagnostic n=153